

Project 215 04 11

050605

POLYBROMINATED DIPHENYL ETHERS (PBDEs) IN SERUM FROM SWEDISH MEN 1988-2002. A LONGITUDINAL STUDY.

Kristina Jakobsson¹, Maria Athanasiadou², Anna Christiansson², Ioannis Athanassiadis², Åke Bergman² and Lars Hagmar¹

¹Department of Occupational and Environmental Medicine, Lund University Hospital, SE-221 85 Lund, Sweden

²Department of Environmental Chemistry, Stockholm University, SE-106 91 Stockholm Sweden

Introduction

Polybrominated diphenyl ethers (PBDEs) have been extensively used as additive flame retardants since the 1970s. Their concentrations have been reported to increase with time both in humans and in wildlife from all environmental compartments¹⁻³. Retrospective time-trend studies conducted using environmental samples originating from the Swedish environment showed a peak in environmental levels in the mid-1980's, after which environmental concentrations decreased or remained unchanged⁴. In Swedish human breast milk, the concentrations of low-medium brominated diphenyl ethers doubled every 5 years until the late 1990s; thereafter a decrease has been indicated, at least for BDE-47⁵. However, there is yet no information on human time trends for hepta-, octa-, nona- and deca-BDEs, neither in Sweden or elsewhere. Further, no longitudinal studies on individual basis have yet been reported.

In 1991 men with a high dietary intake of fish from the Baltic Sea, 12–20 meals/month, had considerably higher BDE-47 levels than men with negligible consumption of fish⁶. These men were resampled in 2001. Time trends for PCB and DDE have been reported elsewhere⁷. Here, we report data on time trends for PBDEs in these men.

Material and methods

Study groups

We had remaining serum (<5 ml) for PBDE-analysis from 37 out of totally 39 men sampled in 1991 and 2001. In 1991 they were between 23 and 69 (median 42) years old. No one was occupationally exposed to PBDEs to our knowledge. Eighteen of these men were stable non-consumers of fish at both sampling

occasions. The remaining nineteen men, who were moderate or high consumers of fatty fish from the Baltic Sea in 1991 (Table 1) had reduced their fatty fish consumption in 2001 compared to ten years earlier; from median 8 (range 4-20) to 3 (1-9) meals per month. For ten of the 37 men serum samples from 1988 and 2002 were also available, remaining from a study of time trends for dioxins and dioxin-like POCs.⁷ Four of these men never ate fish.

Chemical analysis: The chemicals used, extraction of serum, lipid determination, partitioning with an alkaline solution, procedure and analysis have been described in detail elsewhere⁸. Lipids were removed from the extracts by treatment with concentrated sulfuric acid. Fractions containing both the neutral and phenol type substances were subjected to cleanup on two additional steps. First, remaining lipids and endogenous organic material were reduced by running the samples through columns packed with silica mixed with concentrated sulfuric acid (1 g). The analytes were eluted with dichloromethane (10 ml). The last refining cleanup step was performed on a column packed with activated (300°C, 12 h) silica gel (1 g). The silica columns were first prewashed with n-hexane (6 ml) before the samples were applied. A first fraction was collected in n-hexane (3 ml) and a second in dichloromethane (6 ml). The solvent in fraction two was reduced under a gentle stream of nitrogen and replaced with n-hexane prior to gas chromatography-mass spectrometry (GC-MS). Reference compounds, synthesised in house, were used as standards. All solvents were of the highest available commercial grade.

Identification and quantification were performed using a GC-MS Finnigan TSQ 700 (Thermoquest, Bremen, Germany) operating in electron capture chemical ionization (ECNI) mode, tracing the bromide ions (m/z 79 and 81). A DB-5HT

column (15 m × 0.2 mm i.d. and 0.1 µm film thickness) from Supelco (Bellefonte, USA) was used with temperature program of 80°C (1 min) – 15°C/min – 300°C – 2 °C/min – 320°C (2 min). On-column injections were performed using a septum equipped programmable injector fitted with a high performance insert. The injector temperature was 60°C and increased with 150°C per minute up to 300°C for each injection. Helium was used as carrier gas. The transfer line temperature was 290°C and the temperature in the ion-source was 200°C.

Results and Discussion

Fifteen PBDE congeners listed in elution order from the unpolar GC column ; BDE-28, -47, -100, -99, -154, -153, -128, -183, -197, -203, -196, -208, -207, -206 and -209 were detected and quantified.

In the subset of ten men sampled four times between 1988 and 2002, the time trends for selected PBDE congeners are illustrated in Figure 1. Between 1988 and 1991 there was a significant increase of BDE-47 (p=0.05; Wilcoxon signed rank test), followed by a decrease (p=0.12; Friedman test). In contrast, BDE-153 clearly increased between 1991 and 2001, but not thereafter. There was also a tendency of increasing levels of BDE-207 (p=0.11; Friedman test) in these ten men while no tendency was confirmed for BDE-209.

The influence of fish consumption on serum levels of PBDEs in 1991 and 2001 was assessed for the 18 non-consumers and 19 fish-consumers (Table 1; data given for no, moderate and high consumption of fatty fish as determined in 1991). Among the 18 stable non-consumers of fish, the sum of PBDEs increased markedly between 1991 and 2001 (Table 1). The median level increase for low

brominated (3-5 Br) BDEs was 50%, for medium brominated (6-7 Br) BDEs 70%, and for high brominated (8-10 Br) BDEs 90% (the latter not statistically significant).

In the nineteen fish consumers the sum of BDEs did not change over time. The congener pattern of change was different from the non-consumers, with a 40% decrease of low brominated BDEs, 70% increase of medium brominated BDEs, and no significant change in high brominated BDEs. It has to be remembered that these men had reduced their consumption of fatty fish from the Baltic Sea between 1991 and 2001. Thus, the reduced levels of low brominated BDEs may depend on changes in fish consumption habits, rather than changes in environmental levels.

The serum levels of BDE-28, BDE-47, BDE-100, and BDE-154 showed moderate positive correlations (correlation coefficients around 0.3; $p < 0.1$) with fish consumption (measured as meals of fatty fish from the Baltic Sea or as total fish meals per month) in 1991. In samples from 2001 no correlations with these PBDE congener levels were evident, neither for fish consumption data in 1991 nor in 2001. BDE-153 and the hepta- to deca BDEs were not correlated with fish consumption at any point of time.

The men included in our study are not representative of the general Swedish population. They were originally selected in order to represent a wide range of consumption pattern of fatty fish from the Baltic Sea, for studies of the change over time of levels of various persistent organochlorine compounds (POCs). In Sweden, fatty fish from the Baltic Sea has been an important dietary source of these compounds⁹. For PBDEs, our data indicate that the relative contribution

from fatty fish, compared to other sources, may differ between the PBDE congeners, and also over time. Not only diet, but also other sources like air-borne exposure in the home environment are believed to be important¹⁰.

In these men, the average decline of POCs like CB-153, p,p'-DDE and hexachlorobenzene between 1991 and 2001 was 30-50%⁷. In contrast, we observed no over-all decline of the sum of all PBDEs, but marked changes in congener patterns between the two early sampling points and the latter two. This is in accordance with data we have obtained also in some other recent studies¹¹. It is notable that there was a three-fold increase of BDE-153 between 1991 and 2001, regardless of fish consumption habits. In the early 2000s BDE-153 clearly was the dominating PBDE congener. Also, it may be pointed out that the concentrations of BDE-209 were similar to BDE-47 in the samples from the present decade.

Acknowledgements

We gratefully acknowledge the cooperation of Anita Nilsson, RN, Inger Bensryd, RN, Kerstin Kronholm-Diab, RN, and Ewa Wallin, RN for blood sampling. Zoli Mikoczy assisted in the statistical analysis. The study was funded by the Swedish Environmental Protection Agency and the Faculty of Medicine, Lund University, Sweden.

References

- (1) Hites, R. A. *Environ. Sci. Technol.* 2004, 38, 945-956.
- (2) Law, R. J., Alaei, M., Allchin, C. R., Boon, J. P., Lebeuf, M., Lepom, P. and Stern, G. A. *Environ. Int.* 2003, 29, 757-770.
- (3) de Wit, C. *Chemosphere* 2002, 46, 583-624.
- (4) Sellström, U., Bignert, A., Kierkegaard, A., Häggberg, L., de Wit, C. A., Olsson, M. and Jansson, B. *Environ. Sci. Technol.* 2003, 37, 5496-5501.
- (5) Meironytė Guvenius, D. and Norén, K. 303-305. 2001. Stockholm, Sweden. The Second International Workshop on Brominated Flame Retardants. BFR 2001.
- (6) Sjödin, A., Hagmar, L., Klasson Wehler, E., Björk, J. and Bergman, Å. *Environ. Health Perspect.* 2000, 108, 1035-1041.
- (7) Hagmar, L., Wallin, E., Tysklind, M., Sjöström, A., Vessby, B., Jönsson, B. A. G. and Rylander, L. Tidstrender för halter av persistenta klororganiska miljögifter i blod hos vuxna svenska män i relation till konsumtion av fet östersjöfisk. Report to the Swedish Environmental Protection Agency 2004-03-18.
- (8) Hovander, L., Athanasiadou, M., Asplund, L., Jensen, S. and Klasson Wehler, E. *J. Anal. Toxicol.* 2000, 24, 696-703.
- (9) Svensson, B. G., Nilsson, A., Hansson, M., Rappe, C., Åkesson, B. and Skerfving, S. *N. Engl. J. Med.* 1991, 324, 8-12.
- (10) Stapleton, H. M., Schantz, M. and Wise, S. 49-51, Toronto, Canada. The Third International Workshop on Brominated Flame Retardants, BFR 2004.
- (11) Thuresson, K., Bergman, Å. and Jakobsson, K. *Environ. Sci. Technol.* 2005, 39, 1980-1986.




Figure 1. Serum concentrations of BDE-47 (upper left), BDE-153 (upper right), BDE-207 (lower left), and BDE-209 (lower right) (pmol/g lipid weight) in ten men, sampled in 1988, 1991, 2001, and 2002 are presented. Median, 25, 75 percentiles, range and outliers if present are denoted.

Table 1. Concentrations of selected PBDEs (pmol/g lipids, median and range) and sum of PBDEs in serum sampled 1991 and 2001 from a group of Swedish men with varying consumption of fatty fish from the Baltic Sea.

GROUP		BDE-47	BDE-99	BDE-154	BDE-153	BDE-183	BDE-197	BDE-207	BDE-209	3-5 BrBDE ¹	6-7 Br BDE ²	8-10 Br BDE ³	sumBDE ⁴
All subjects, (n=37)		2.1	0.40	1.3	0.93	0.41	0.45	0.61	1.3	3.5	2.8	3.0	11
	1991	0.35-17	0.16-7.4	0.15-5.6	0.34-3.7	0.11-5.4	0.05-14	0.07-7.7	0.01-21	0.68-30	0.64-11	0.25-35	3.3-59
		1.6	0.64	1.1	3.0	0.46	0.58	0.70	1.4	3.1	4.8	3.0	14
	2001	0.36-13	0.13-6.3	0.30-3.2	0.66-34	0.17-1.9	0.18-8.3	0.31-2.8	0.47-5.2	0.66-18	1.2-36	1.5-16	4.2-57
	p-value ⁵	p=0.09	p=0.04	-	p<0.001	-	p=0.19	-	-	-	p<0.001	-	-
Fish consumption ⁶													
None (n=18)		1.2	0.31	1.3	0.78	0.39	0.54	0.42	0.39	1.6	2.5	1.7	6.0
	1991	0.35-8.1	0.16-2.5	0.31-2.4	0.40-3.7	0.11-3.8	0.05-14	0.07-7.7	0.01-9.4	0.68-13	1.0-8.9	0.25-35	3.3-47
		1.5	0.73	1.1	2.5	0.52	0.61	0.69	1.5	2.4	4.2	3.2	13
	2001	0.54-5.3	0.30-2.5	0.40-2.9	1.2-8.6	0.27-1.9	0.34-8.3	0.31-2.8	0.47-3.4	1.0-9.5	2.1-10	1.5-16	6.2-26
	p-value ⁵	-	p=0.003	-	p<0.001	-	-	-	-	p=0.09	p=0.001	-	p=0.03
Moderate (n=10)		4.7	0.88	1.3	1.6	0.46	0.65	1.2	2.7	7.7	3.4	4.7	17
	1991	0.77-17	0.17-7.4	0.53-3.1	0.60-2.8	0.14-5.4	0.16-1.3	0.09-2.1	0.01-21	1.3-30	1.8-11	0.38-27	8.6-59
		2.6	0.91	1.4	3.9	0.40	0.60	0.76	1.1	4.7	5.7	2.8	14
	2001	0.36-13	0.13-3.0	0.30-3.2	0.66-9.4	0.25-1.6	0.24-0.87	0.35-1.0	0.75-2.5	0.66-18	1.2-14	2.1-5.3	4.2-27
	p-value ⁵	p=0.08	-	-	p=0.007	-	-	p=0.04	p=0.17	p=0.09	p=0.09	-	-
High (n=9)		3.4	0.51	1.4	1.0	0.42	0.29	0.44	0.50	5.1	2.9	1.8	12
	1991	1.2-7.2	0.32-1.2	0.15-5.8	0.34-1.7	0.14-3.5	0.15-0.68	0.17-1.2	0.01-3.2	2.3-13	0.6-11	0.21-5.8	5.9-22
		1.4	0.49	1.3	3.1	0.33	0.47	0.82	1.3	2.5	5.0	3.2	15
	2001	0.99-7.6	0.37-6.3	0.74-3.0	1.7-34	0.17-0.58	0.18-0.88	0.35-2.3	0.63-5.2	2.0-16	2.8-36	1.7-11	7.5-57
	p-value ⁵	p=0.11	-	-	p=0.008	-	p=0.09	-	-	p=0.11	p=0.07	-	-

¹sum of BDE-28,-47, -100, -99.

²sum of BDE--154, -153, -128,-183

³sum of BDE-197, -203, -196, -208, -207, -206, -209.

⁴sum of BDE-28,-47, -100, -99, -154, -153, -128,-183, -197, -203, -196, -208, -207, -206, -209.

⁵Wilcoxon paired test. p-levels >0.2 are not given in the table

⁶Consumption of fatty fish from the Baltic sea as classified in 1991. Moderate: 4-8 meals/month. High: 12-20 meals/month.